(D)

OFFICE OF NAVAL RESEARCH

AD-A256 408

END-OF-YEAR REPORT

PUELICATIONS/PATENTS/PRESENTATIONS/HONORS/STUDENTS REPORT

for

GRANT or CONTRACT: N00014-91-J-1617

R&T Code 4132054--01

Technical Report No. 1



Synthesis of a Low Dielectric Perfluoromethylene Linked Cyanate Resin

by

Arthur W. Snow, James R. Griffith, Robert L. Soulen, Jeffery A. Greathouse, J.K. Lodge

Chemistry Department
Southwestern University
Georgetown, Texas
78626
April 9, 1992

Reproduction in whole or in part is permitted for any purpose of the United States Government.

This document has been approved for public release and sale: its distribution is unlimited.



REPORT DOCUMENTATION PAGE					Form Approved OMB No 0704 0188		
1. REPORT SECURITY CLASSIFICATION		16 RESTRICTIVE MARKINGS					
Unclassified							
2. SECURITY CLASSIFICATION AUTHORITY		3 DISTRIBUTION/AVAILABILITY OF REPORT					
26 DECLASSIFICATION DOWNGRADING SCHEDU	.E	Approved for public release Distribution Unlimited					
4 PERFORMING ORGANIZATION REPORT NUMBER	R(S)	5 MONITORING ORGANIZATION REPORT NUMBER(S)					
Technical Report #1							
6a NAME OF PERFORMING ORGANIZATION 66 OFFICE SYMB		78 NAME OF MONITORING ORGANIZATION					
Southwestern University (If applicable)		ONR					
5c. ADDRESS (City, State, and ZIP Code)		7b ADDRESS (City, State, and ZIP Code)					
Chemistry Department		Department of the Navy					
University at Maple		Arlington, VA 22217					
Georgetown, Texas 78626							
8a NAME OF FUNDING / SPONSORING ORGANIZATION	8b OFFICE SYMBOL (If applicable)	9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER					
ONR 8c ADDRESS (City, State, and ZIP Code)			TO STATE AND ADD DO				
		10 SOURCE OF FUNDING NUMBERS PROGRAM PROJECT TASK WORK UNIT			WORK UNIT		
Office of the Navy		ELEMENT NO	NO	NO	ACCESSION NO		
Arlington, VA 22217		N00014-91	J-1617	ŧ			
11 TITLE (Include Security Classification) Synthese of a Low Dielectric Pe	erfluoromethyle	ne Linked Cy	yanate Resi	n			
12 PERSONAL AUTHOR(S)				. ,			
Arthur Snow, James Griffith, Je	attery Greathou	ise. Robert S 14 Date of Repor	Soulen, Jay	Lodge	PAGE COUNT		
FROM	10	9274/9	m (rear, worth, r	20,7	2		
16 Supplementary Notation Paper published in the proceed and Engineering, San Francisco.	, 1992.						
17 COSATI CODES	18 SUBJECT TERMS (C			_			
FIELD GROUP SUB GROUP	low dielectri		cyanate				
	fluoropolymer		cyanura	te iii	K		
19 ABSTRACT (Continue on reverse if necessary and identify by block number) Copper coupling of 1,6-diiodoperfluorohexane and para iodoanisole gave 1,5-bis (4'-methoxyphenyl) perfluorohexane in good yield. Demethylation with boron tribromide and subsequent reaction with cyanogen bromide gave 1,5-bis (4'-cyanatophenyl) perfluorohexane. Polymerization of the cyanate with copper acetylacetonate provided a good exotherm for the desired trimerization reaction. Measured dielectric permittivity between 2.3 and 2.4. Discussion of attempted preparation of the free phenol perfluoroalkane linked compounds is presented.							
20 DISTRIBUTION AVAILABILITY OF ABSTRACT	21 ABSTRACT SECURITY CLASSIFICATION						
UNCLASSIFIED UNUMITED SAME AS F	RPT DTIC USERS	226 TELEPHONE (/	In the American	1122. 60	tict (VAADO)		
559 MV-AIS OF REPEGINDING HIDINIDIDA		226 TELEPHONE (I	include Area Code,	1/22 6	FILE STIVIBUL		

SECURITY CLASSIFICATION OF THIS PAGE
•



POLYMERIC MATERIALS SCIENCE AND ENGINEERING



VOLUME 66

	j
tosossian For	
TOPS CASAL ONE THAT SHOULD SEE COMME	<u>n</u>
	Undes
A-1	2/or 1

SPRING MEETING 1992 SAN FRANCISCO, CALIFORNIA

PROCEEDINGS OF THE AMERICAN CHEMICAL SOCIETY DIVISION OF POLYMERIC MATERIALS: SCIENCE AND ENGINEERING

• 1992 American Chemical Society

SYNTHESIS OF A LOW DIELECTRIC PERFLUOROHEXAMETHYLENE LINKED CYANATE RESIN

A.W. Snow and J.R. Griffith Naval Research Laboratory Washington, D.C. 20375

R.L. Soulen and J.A. Greathouse Chemistry Department, Southwestern University Georgetown, Texas 78626

> J.K. Lodge SFA Inc. Landover, Maryland '20785

INTRODUCTION

The objective of this work is to synthesize a polymer network wherein the chain structure is polytetrafluoroethylene and the network junction is a triazine linkage. The motivation is to achieve a resin with the dielectric properties of Teflon and the mechanical, thermal and processing properties of a good thermoset. With respect to application as a printed wiring board laminating resin this structure appeared to possess the features of high symmetry, low polar content and low polarizibility symmetry, low point content and point of proscribed as desirable for low dielectric thermoset resins [1]. Inspection of the above described structure suggests that a nitrile trimerization of an α, ω -perfluoromethylenedinitrile is an obvious synthetic route. However, nitrile trimerization requires severe temperature conditions and is not a high yield reaction. Interestingly, a successful route to this perfluoromethylenetriazine network structure via a perfluoroamidine monomer has been reported 31 years ago (2), but it involved reaction temperatures of 300 to 400°C and condensation of ammonia.

The cyanate trimerization appeared to confer processibility at reasonable cure temperatures with minimal sacrifice of the symmetry and low polarity characteristics of the above mentioned structure. The monomer and polymer we identified for synthesis are depicted in the following polymerization reaction.

In the synthesis and trimerization of cyanates, fluorocarbon chemistry has been found to work a little differently from hydrocarbon chemistry [3]. In this respect, the fluoroalkylphenylcyanate depicted above was similar in reactivity to hydrocarbon phenylcyanates, but different in synthesis due to an instablilty of the fluoroalkylphenol. In this paper we present the synthesis and polymer-ization and characterization of 1,6-bis(4'-cyanatophenyl)dodecafluorohexane (BP12FCY).

EXPERIMENTAL

All reagents and solvents were of reagent-grade quality, purchased commercially and used without further purification unless otherwise noted. Measurements and associated instrumentation are: infrared spectra, Perkin-Elmer Model 1320; 'H and 'FT NNR, Varian Models 360 and 390 using TMS and CFC13 as internal standards; mass spectra, Varian Saturn Model GC/MS; DSC, DuPont 2100 Thermal Analysis System with a 910 differential scanning calorimeter, permittivity, HP8510 Network Analyzer (0.5-18 GHz). All melting points are uncorrected. 1,6-Pis(4'-methoxyphenyl)dodecafluorohexane(BP)2FCMe). A mixture of 18.76 g (80.1 mmol) of 4-iodoanisole, 1:11 g (20.0 mmol) of 1,6-diodododecafluorohexane and 6.35 g (100 mmol) of copper powder in 5 g DMSO was reacted at 125-135°C (24 hr) then 165°C (1.5 hr) followed by sublimation at 110°C/0.025 mmHg. Resublimation gave 8.5 g product as white

crystals: mp 77.0-78.5°C; H NMR (CDC13) 3.80 ppm s (3H), 6.93 ppm d (2H) and 7.50 ppm d (2H); F NMR (CDC13) -107.2 ppm (ATCF₂-) and 119.3 ppm (-(CF₂)₂-); IR see figure 1.

1.6-Bis(4'-cyanatophenylidodecafluorohexane (BP12FCY).

By gas tight syringe 1.91 ml (5.04 g, 20.2 mzol) of boron tribromide were added to 40 ml freshly distilled methylene chloride under nitrogen followed by cooling to 0-5°C and dropwise addition of 4.51 g (8.77 mmol) of BP12FCMH dissolved in 100 ml of freshly distilled CH₂Cl₂. After stirring 12 hr at 20°C, the reaction mixture was poured into 150 ml rapidly stirred water, and the organic layer was repeatedly extracted with 200 ml portions water until the water extract was neutral. The CH₂Cl₂ solution was dried over Na₂SO₄, filtered, and IR spectrum of evaporated film checked for methoxy cleavage. To this solution were then added 1.86 g (17.5 mmol) of cyanogen bromide followed by cooling to 5°C (ppt formation at lower temperatures) and dropwise addition of 1.77 g (17.5 mmol) triethylamine dissolved in 6 ml of CH₂Cl₂. The reaction mixture was stirred at 0-5°C for 2.5 hr then allowed to slowly to warm to room temperature with stirring for an additional hour. The reaction mixture was worked up by pouring into 500 ml of stirred water, two extractions with 200 ml of water, Na₂SO₄ drying and evaporation. A tedious recrystalization from cyclohexane yielded 1.70 g (39%) of BP12FCY: mp 75-77°C; H NOR (CDC1) 3.80 ppm (residual -OCH3) and 6.8-7.7 ppm (Ar-B); F NMR (CDC1) -158 ppm (4F) and -169 ppm (8F); IR see figure 1; MS no molecular ion, 143 (100%) CF₂CH₃OH.

BP12FCY Cure. A mixture of 0.990 g BP12FCY and 0.0089 g catalyst solution (0.30 g copper acetylacetonate in 4.00 g nonylphenol [4] was degassed under vacuum (~1 mmHg)/100°C, mixed by stirring after melting and cured at 115°C for 1.5 hr followed by 180°C for 1 hr.

RESULTS AND DISCUSSION

The synthesis of BP12FCY was undertaken by the route depicted in figure 1. The initial plan was to couple p-iodophenol with the diiodoperfluoro-However, much difficulty was experienced in isolating a product, and it appears that the pfluoromethylene substituted phenol is not stable to purification operations. Protecting the phenol with the methoxy group by starting with p-fluoroanisole in place of p-fluorophenol solved this problem, and the bis(methoxyphenyl)perfluorohexane could be separated from the DMSO and purified by sublimation.

The problem now became one of converting the methoxy groups to cyanate groups without having the intermediate phenol undergo decomposition. was needed is a mild method of cleavage of the anisole and preferably a medium which would allow proceeding immediately to the cyanation reaction without isolating the phenol. Typically, cleavage by hydroiodic or sulfonic acids require tempera-tures in excess of 100°C [5], however, boron tribromide is reported to effectively demethylate methyl phenyl ethers at or below room temperature in methylene chloride solution [6]. This was particularly attractive because the methylene chloride medium could easily be made neutral and is a good solvent for the low temperature base catalyzed cyanation reaction. An excess of BBr; was used, and it appeared from the hydroxyl absorption in the infrared spectrum (figure 2) that a good conversion had been obtained although a residual methoxy is also present.

The cyanation reaction was undertaken after obtaining a neutral aqueous extract and drying the methylene chloride solution. The infrared spectrum (figure 2) indicated that the hydroxyl was quantitatively converted to the cyanate group (2250 cm⁻¹).

Proceeding to the polymerization uncatalyzed cure of the BP12FCY monomer was first attempted. In this instance the DSC thermogram displayed a broad exotherm with a maxima at 297°C which we attributed to the trimerization reaction and a second exotherm onset at ~350°C which we suspect may be associated with decomposition. It has been demonstrated that addition of catalytic quantities of transition metals (copper in particular) in combination with a phenol can markedly lower the

Figure 1. Synthesis of 1,6-Bis(4'-cyanatonhenyl)perfluorohexane.




Figure 2. Infrared Spectra of Perfluorohexanedicyanate, Precursors, Monomer and Resin.

cure temperature and provide much better control over the trimerization exotherm [4]. Addition of 0.89 phr of a copper acetylacetonate/nonylphenol catalyst markedly narrowed and lowered the trimerization exotherm to a 167°C maximum in the DSC thermogram (figure 3). Following a curing schedule of 115°C for 1.5 hr and 180°C for 1 hr, an infrared spectrum indicated the cyanate (2250 cm⁻¹) was completely consumed and a large amount of triazine (1573 and 1375 cm⁻¹ [7]) had formed (figure 2).

A measurement of the dielectric permittivity presented in figure 4. Its response to Its response to




Figure 3. DSC Thermogram of BP12FCY Cure Catalyzed With 0.89 phr CuAcAc/Nonylphenol.




Figure 4. Dielectric Measurement on BP12FCY

Dielectric Permittivity of Thermoset and Thermoplastic Resins [1,8].

Resin	Permittivity
FR-4 epoxy	3.6
Polyimide (BMI)	3.2
Polyetherimide	3.1
Polysulfone	3.0
Polycarbonate	3.0
Polybutadiene	2.8
Polyolefin	2.3
Polytetrafluoroethylene	2.0
Cyanates	2.0
**<0∙ (}+(}`oc*	3.1
40€ >+ DD 12000	
~~ ~~ ~~ ~~ ~~ ~~ ~~ ~~ ~~ ~~ ~~ ~~ ~~	2.8
MCO Con Coon	2.75
n _y /	2.73
roAlAon	2.66
W & W	

frequency is relatively flat with a value between 2.3 and 2.4. Compared with other cyanates in specific [8] and thermosets or thermoplastics in general [1] this value appears reasonable for the polymer structure. Table 1 presents permittivity values with various associated resins and structures but has ignored any dependence on frequency. The BP12FCY result presented here is intermediate to the 2.0 for PTFE and the 2.66 for the dicyanate of bisphenol F. The true BP12FCY result may be slightly lower then that reported here since there is a residual quantity of methoxy groups in this synthesis as evidenced by the infrared band at 2845 \mbox{cm}^{-1} in figure 2.

to lower the In summary, to lower the dielectric permittivity of a thermoset resin, the approach of connecting perfluoromethylene chains with cyanurate linkages provides a structure that is near equivalent to polyethylene at perfluoromethylene chain length of six. Increased fluoroalkyl content will more closely approach the value of Teflon.

REFERENCES

- S.J. Mumby, J. Elect. Mat. 18, 241 (1989). H.C. Brown, J. Poly. Sci. 44, 9 (1960). A.W. Snow and J.R. Griffith, J. Fluorine Chem.
- A.W. Snow and J.R. Stratter, 15, 471 (1980).
 D.A. Shimp, PMSE Preprints 54, 107 (1986).
 J. March, Advanced Organic Chemistry, 3rd ed.,
 J. Wiley & Sons, New York (1985) pp. 384, 358.
 J.F.W. McOmie, M.L. Watts and D.E. West,
- Tetrahedron 24, 2292 (1968). R.F. Cozzens, P. Walter and A.W. Snow, J.

- Appl. Poly. Sci. 34, 601 (1987). D.A. Shimp, J.R. Christenson and S.J. Ising, 34th Int. SAMPE Symp. Proc., pp 222-233, May 8-11, 1989.

FINANCIAL STATUS REPORT

To Which Report is Submitted	By Federal Agency No. 0348-0039			pages	
AVY	N00014-91-J-1617				
. Recipient Organization (Name and complete add	ress, including ZIP code)	L		·	
Southwestern University, l	University at Mapl	e, Georgetown,	TX 78626		
RA414	5. Recipient Account Number or Identifying Numbe 6. Final Report (X.) Yes () No			7. Basis (X) Cash () Accrual	
 Funding/Grant Period (See Instructions) From (Month, Day, Year) 4/1/91 	To: (Month, Day, Year) 12/31/91		9. Period Covered by this Report From: (Month Day, Year) 4/1/91		
10. Transactions:		Previously Reponed	II This Period	III Cumulative	
Total Outlays		0 00	\$10,000.00	\$10,000.00	
b. Recipient share of outlays		0 00	0	0	
c. Federal share of outlays	·····	0.00	\$10,000.00	\$10,000.00	
d. Total unliquidated obligations					
Reciplent share of unliquidated obligations					
f. Federal share of unliquidated obligations					
g. Total Federal share (Sum of lines c and 1)					
h. Total Federal funds authorized for this fund	Ing period				
1. Unobligated balance of Federal funds (Line h	mínus line g)				
a. Type of Rate (Place 'X' in () Provisional (X) Predetermin					
11, Indirect b Rate c Base	d. Total Amou e. Federal	Share			
Expense N/A N/A 12. Remarks: Attach any explanations deemed legislation.	N/A N/A necessary or Information requ	ulted by Federal sponsori	ng agency in comp	liance with governing	
13. Certification: I certify to the best of my k unliquidated obligations are				hat all outleys	
Type or Printed Name and Title Richard L. Anderson,	Telephone (Area c	Telephone (Area code, number and extension) (512) 863-1475			
Signature of Authorized Certifying Official	Ì	Date Report Submitted June 1, 1992			

OFFICE OF NA L RESEARCH PUBLICATIONS/PATENTS/PRESENTATIONS/HONORS/STUDENTS REPORT

1. Other funding: Robert A. Welch Foundation-Departmental Grant \$8000/91-92; \$24,000/90-92

OFFICE OF NAVAL RESEARCH

PUBLICATIONS/PATENTS/PRESENTATIONS/HONORS REPORT

Supplemental page R&T Number: 4132054__01 Grant Number: N0014-91-J-1617

Item d:

Printed technical reports & non-refereed papers:

- 1) A.W. Snow, J.R. Griffith, R.L. Soulen, J.A. Greathouse and J.K. Lodge; Synthesis of a Low Dielectric Perfluoromethylene Linked Cyanate Resin; Proceedings of the ACS Division of Polymeric Materials Sciences and Engineering, San Francisco, 1992
- 2) Jeffery A. Greathouse; A Method of Synthesizing Perfluoroalkyl-substituted Aromatic Compounds; April 8, 1992, Senior Research, Southwestern University
- 3) Audrey Lim; A Study of the Water Permeability of Fluoroacrylate Films as Compared to Commercial Films; April 27, 1992, Senior Research, Southwestern University

Item j;

Honors

William Currington Finch Award to Robert L. Soulen, May 9, 1992

END

DATE: //-92

DTIC